

Bactroban® Nasal Ointment (Mupirocin)

FORM Mupirocin 2% Nasal Ointment

INDICATION Eradication of nasal staphylococci, including methicillin-resistant Staphylococcus aureus (MRSA).

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
All ages	One Application	2-3 times daily for 5-7 days	Apply to the inside of each nostril

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

FURTHER INFORMATION Bacterial resistance has emerged, particularly during long term treatment

LICENSED STATUS Licensed for use in all ages

LINKS [BNF for Children](#) / [Electronic Medicines Compendium](#)
APPLICABLE POLICIES [West of Scotland Neonatal Guidelines](#):

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo pharm group	Checked by	Michael Mitchell
Date prepared	Aug 2016	Date updated	October 2021
Updated by	Lauren Williams	Review Date	October 2024

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications. This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.