

## Oxybutynin

**FORM** Liquid 2.5mg/5ml  
**INDICATION** Neurogenic bladder instability

### DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate to six months	100 – 200 micrograms/kg	2 -3 times daily	Oral

Maximum dose – 1.25mg THREE times daily

### CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

### FURTHER INFORMATION

### STORAGE

**LICENSED STATUS** Unlicensed dosage

**LINKS** [BNF for Children:](#) / [Electronic Medicines Compendium](#)

**APPLICABLE POLICIES** [West of Scotland Neonatal Guidelines:](#)

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	Peter Mulholland	Checked by	WoS Neonatal Pharmacists
Date prepared	January 2017	Date updated	October 2021
Updated by	Peter Mulholland	Review Date	October 2024

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications. This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.